

Far Eastern Entomologist

Дальневосточный энтомолог

Journal published by Far East Branch
of the Russian Entomological Society
and Laboratory of Entomology,
Institute of Biology and Soil Science,
Vladivostok

Number 323: 1-6

ISSN 1026-051X

October 2016

<http://urn:lsid:zoobank.org:pub:9AFFBC50-55E4-4A9C-A7F6-E3EC9AD31C01>

A NEW SPECIES OF THE GENUS *DICROTENDIPES* KIEFFER, 1913 (DIPTERA: CHIRONOMIDAE: CHIRONOMINAE) FROM NORTH KOREA

O. V. Orel*, E. A. Makarchenko

*Institute of Biology and Soil Science, Far Eastern Branch of the Russian
Academy of Sciences, Vladivostok 690022, Russia. *Corresponding author, E-mail:
zorina@biosoil.ru*

Dicrotendipes koreanus Orel, **sp. n.** is described on the base of adult male from
North Korea. New species differs from congeners in the structure of hypopygium.

KEY WORDS: Diptera, Chironomidae, *Dicrotendipes*, new species, Korea.

О. В. Орел*, Е. А. Макаrenchенко. Новый вид рода *Dicrotendipes* Kieffer,
1913 (Diptera: Chironomidae: Chironominae) из Северной Кореи // Дальне-
восточный энтомолог. 2016. N 323. С. 1-6.

Из Северной Кореи по самцу описан новый для науки вид комаров-звонцов
Dicrotendipes koreanus Orel, **sp. n.** Новый вид отличается от других видов рода
строением гипопигия.

Биолого-почвенный институт ДВО РАН, Владивосток 690022, Россия.
*Корреспондирующий автор.

INTRODUCTION

The genus *Dicrotendipes* was erected by Kieffer in 1913, with *Dicrotendipes
septemmaculatus* (Becker, 1908) as a type species (Kieffer, 1913). The immature

stages of *Dicrotendipes* are found in both lentic and lotic habitats, but are generally more prevalent in lentic conditions (Epler, 1988). More than 100 species have been described (Qi *et al.*, 2012). Twenty three species of *Dicrotendipes* have been recorded from the Eastern Palaearctic region, namely, 10 species from Japan (Niitsuma, 1995; Sasa & Kikuchi, 1995), 10 species from China (Qi *et al.*, 2016), 10 species from Russian Far East (Makarchenko *et al.*, 2005; Zorina, 2006), and 6 species from Korea; beside them five species were known from South Korea, namely, *D. hannervosus* Ree, 2013, *D. nervosus* (Staeger, 1839), *D. pelochloris* (Kieffer, 1912), *D. septemmaculatus* (Becker, 1908), *D. tamaviridis* Sasa, 1981, and two species, *D. nervosus* and *D. septemmaculatus*, from North Korea (Riess, 1980; Ree, 1981; Ree & Kim, 1981; Ree *et al.*, 1995; Ree, 2013; Ree, 2014). One new species of *Dicrotendipes* is found in North Korea and described below.

Present paper is based on the material collected in North Korea by the Polish professor Wiesław Krzemiński in 1981. Material was fixed in 70% ethanol and mounted in For-Berlese solution.

Morphological terminology and abbreviations follow Sæther (1980) and Epler (1988).

Holotype of new species is deposited in the National Institute of Biological Resources, Incheon, Republic of Korea (NIBR).

DESCRIPTION

Family Chironomidae

Subfamily Chironominae

Dicrotendipes koreanus Orel, sp. n.

Figs 1–8


TYPE MATERIAL. Holotype – male, **Democratic People's Republic of Korea**: Pyongyang (= Phjôngjang or Phenian), small lake, 11.VI 1981, coll. W. Krzemiński (NIBR).

DESCRIPTION. ADULT MALE (n=1). General colour yellowish brown. Total length 6.0 mm; total length / wing length 2.2.


Head. Frontal tubercles rudimentary (3,5 µm Ø circles free of microtrichia). Temporal setae 15. Clypeus with 24 setae. Maxillary palp 600 µm, lengths of last 4 palpal segments (µm): 72, 200, 144, 184. Palp length / head width 0.70. Scapus yellowish, flagellomeres 1–11 – dark-brown, 1764 µm long. AR 3.0. Antenna length / palp length 2.94.

Thorax. Ground colour of scutum pale yellow, mesonotal strips yellowish brown. Aps 0, Ac 22, Dc 18–21, Pa 8, Su 2. Scutellum pale yellow, with 18 setae. Postnotum in distal 2/3 yellowish brown.

Wing length 2.7 mm. Squama with 15–16 setae; brachiolum with 2–3 setae. R, R₁ with 43–52 setae, R₄₊₅ with 22–29 setae. VR 1.11. Halter's pale yellow.



Figs. 1–4. Details of the hypopygium structure of *Dicrotendipes koreanus* sp. n. 1 – hypopygium, dorsal view; 2 – tergite IX, lateral view; 3 – superior and inferior volsellae, lateral view; superior and inferior volsellae, dorsal view. Scale bar 50 μ m.



Figs. 5–8. Hypopygium of *Dicrotendipes koreanus* sp. n. 5 – dorsal view of hypopygium; 6 – ventral view of hypopygium; 7 – anal point and superior volsella, dorsal view; 8 – anal point and superior volsella, ventral view. Scale bar 50 μ m.

Legs yellowish brown. Terminal combs of t_3 with spurs long 41 μm and 27 μm (spurs on t_2 broken). BR_2 3.13. Mid ta_1 sensilla chaetica 17–20. Length and proportions of legs segments are as follow:

P	f	t	ta_1	ta_2	ta_3	ta_4	ta_5	LR	SV	BV
P_1	1260	1113	—	—	—	—	—	—	—	—
P_2	1155	1155	630	420	294	147	126	0.55	3.67	2.98
P_3	1407	1512	—	—	—	—	—	—	—	—

Abdomen yellowish brown.

Hypopygium (Figs 1–8). Tergite IX with 12–14 median setae set within two clearly defined ovoid areas. Laterosternite IX with 6–7 setae. Anal point short and wide (length 85 μm , width 75 μm), with 16 lateroventral setae on each side; in lateral view hunchbacked and straight. Gonocoxite 286 μm long, with 6–7 setae. Superior volsella semi-circular shape (length 109–119 μm), slightly curved, and the most extended in the apical third (44 μm), microtrichiose, and with 7–8 strong ventromedial setae; LWR 2.5–2.7. Inferior volsella (170 μm long) apically clubbed and curved in lateral view, and covered with microtrichia except of apical part, with 10 long dorsal and 8–9 short ventral setae. Gonostylus 211 μm long, massive and straight, expanded in the proximal two-third (61 μm), gradually narrows towards the apex. HR 1.36.

COMPARISION. Adult male of *D. koreanus* sp. n. clearly differs from all known species of *Dicrotendipes* by the original structure hypopygium as well as by the following combination of features: yellowish brown colour; total length 6.0 mm; frontal tubercles small; AR 3.0; anal point short and wide, in lateral view hunchbacked and straight; superior volsella semi-circular shape slightly curved, microtrichiose, with 7–8 strong ventral setae; inferior volsella clubbed apically and curved in lateral view; gonostylus massive and straight, expanded in the proximal two-third, gradually narrows towards the apex; HR 1.36.

ETYMOLOGY. Latin noun (*Korean*) and suffix *-us*. The species is named for the country when material was collected.

DISTRIBUTION. Known only from North Korea.

ACKNOWLEDGEMENTS

Author is grateful to Dr. Gilka from Department of Invertebrate Zoology, University of Gdańsk, Poland for the transferred material collected by Dr W. Krzemiński in the Northern Korea. This study was performed in the frames of the state research project No. 01201361586 and was supported by the grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR201601203).

REFERENCES

- Epler, J.H. 1988. Biosystematics of the genus *Dicrotendipes* Kieffer, 1913 (Diptera: Chironomidae: Chironominae) of the world. *Memoirs of American Entomological Society*, 36: 1–214.
- Kieffer, J.J. 1913. Chironomidae and Cecidomyiidae. Voyage de Ch. Alluaud et R. Jeannel en Afrique orientale, 1911–1912. *Résultats scientifiques, Insectes Diptères*, 1–43.
- Qi, X., Lin, X.L. & Wang, X.H. 2012. Review of *Dicrotendipes* Kieffer from China (Diptera: Chironomidae). *ZooKeys*, 183: 23–36.
- Qi, X., Peng, J., Zhang, R. & Wang, X. 2016. *Dicrotendipes weiqiangensis* sp. nov. (Diptera: Chironomidae) from Zhejiang, Chine. *Entomotaxonomia*, 38(3): 1–5.
- Makarchenko, E.A., Makarchenko, M.A., Zorina, O.V. & Sergeeva, I.V. 2005. Preliminary data of fauna and taxonomy of chironomids (Diptera, Chironomidae) of the Russian Far East. *Vladimir Ya. Levanidov's Biennial Memorial Meetings*, 3: 394–420. [In Russian].
- Niitsuma, H. 1995. Three species of the genus *Dicrotendipes* (Diptera, Chironomidae) from Japan. *Japanese Journal of Entomology*, 63(2): 433–449.
- Ree, H. 2013. Six new and two unrecorded species of Chironomidae (Insecta: Diptera) from Korea. *Entomological Research*, 43: 322–329.
- Ree, H. 2014. Nine new to science and eleven unrecorded species of Chironomidae (Diptera) from Korea. *Entomological Research Bulletin*, 30(1): 33–60.
- Ree, H. & Kim, H.S. 1981. Studies on Chironomidae (Diptera) in Korea. 1. Taxonomical study on adults of Chironomidae. *Proceeding of the College of Natural Sciences, Seoul National University*, 6(1): 123–226.
- Ree, H., Lee, W.J., Lee, I.Y. & Jeon, S.H. 1995. Fauna of Arthropods of medical importance in Chindo Island. *Korean Journal of Systematic Zoology*, 11(1): 87–100.
- Riess, F. 1980. Zur Zoogeographie der Chironomidenfauna Nordkoreas. P. 145–149. In: Murray D.A. (Ed.). *Chironomidae. Ecology, systematics, cytology and physiology*, Pergman Press, Oxford and New York.
- Sæther, O.A. 1980. Glossary of chironomid morphology terminology (Chironomidae, Diptera). *Entomologica Scandinavica*, Suppl.14: 1–51.
- Sasa, M. & Kikuchi, M. 1995. *Chironomidae (Diptera) of Japan*. University of Tokyo Press. 333 pp.
- Zorina, O.V. 2006. *Dicrotendipes orientalis* sp. n., a new species of non-biting midges (Diptera: Chironomidae: Chironominae) from the Russian Far East. *Russian Entomological Journal*, 15(1): 79–82.