

TO THE TAXONOMY OF *HARPACTUS VEDICUS* NESTEROV, 1994
(HYMENOPTERA, CRABRONIDAE, BEMBICINAE)

M. V. Mokrousov¹⁾, M. Yu. Proshchalykin^{2*)}, M. M. Maharramov³⁾

1) Institute of Biology and Biomedicine at Lobachevsky State University of Nizhny Novgorod, Nizhny Novgorod, 603950, Russia.

2) Federal Scientific Center of the East Asia Terrestrial Biodiversity, Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, 690022, Russia. *Corresponding author. E-mail: proshchalykin@biosoil.ru

3) Nakhchivan State University, Nakhchivan, AZ 7012, Azerbaijan.

Summary. *Harpactus vedicus* Nesterov, 1994 is resurrected from synonymy of *H. elegans* (Lepeletier de Saint Fargeau, 1832). A new synonymy is proposed for *Harpactus vedicus* Nesterov, 1994 = *H. transcausicus* Nemkov, 1994, **syn. n.**

Key words: Apoidea, Spheciformes, new synonymy, Palaearctic region, Azerbaijan, Nakhchivan Autonomous Republic.

М. В. Мокроусов, М. Ю. Прощалькин, М. М. Маггеррамов. К таксономии *Harpactus vedicus* Nesterov, 1994 (Hymenoptera, Crabronidae, Bembicinae) // Дальневосточный энтомолог. 2024. N 499. С. 19-24.

Резюме. *Harpactus vedicus* Nesterov, 1994 (Hymenoptera, Crabronidae, Bembicinae) восстанавливается из синонима *H. elegans* (Lepeletier de Saint Fargeau, 1832) и рассматривается как валидный таксон. Предложена новая синонимия: *Harpactus vedicus* Nesterov, 1994 = *H. transcausicus* Nemkov, 1994, **syn. n.**

INTRODUCTION

Digger wasps of the genus *Harpactus* Shuckard, 1837 are represented by 73 species occurring in Afrotropical, Holarctic and Oriental zoogeographical regions. Six species are known from the Nakhchivan Autonomous Republic, Azerbaijan (Mokrousov *et al.*, 2019). Morphologically, the genus is well separated in the subtribe Gorytina (Bohart & Menke, 1974; Nemkov & Lelej, 2013), but the species require revision.

MATERIAL AND METHODS

Studied material included type series (2 ♀, 12 ♂) of *Harpactus transcausicus*, 15 specimens of *H. elegans* and 97 specimens of *H. vedicus* collected in 2019–2022 in the Nakhchivan Autonomous Republic (Azerbaijan) by M. Yu. Proshchalykin, Kh.A. Aliyev and

M.M. Maharramov. Specimens were deposited in the collection of the Federal Scientific Center of the East Asia Terrestrial Biodiversity of the Far Eastern Branch of the Russian Academy of Sciences, Vladivostok, Russia [FSCV], the personal collection of M.V. Mokrousov, Nizhny Novgorod, Russia [MMC] and the Zoological Institute of the Russian Academy of Sciences (Russia, St. Petersburg) [ZISP].

Photographs were taken using a Canon M200 digital camera attached to an Olympus SZX16 stereomicroscope. Multifocus images were created from stacks of photographs using Helicon Focus v. 7.7.4 Pro software. The final images were processed for sharpness, contrast and brightness using Adobe® Photoshop® software.

The classification and distribution generally follows Pulawski (2024).

TAXONOMY

Genus *Harpactus* Shuckard, 1837

Arpactus Jurine, 1807: 192, junior homonym of *Arpactus* Panzer, 1805, and of *Arpactus* Panzer, 1806. Type species: *Arpactus formosus* Jurine, 1807, designated by Shuckard, 1837: 220.

Harpactus Shuckard, 1837: 221. Emendation of *Arpactus* Jurine, 1807 (see Pulawski, 1985: 59).

Harpactes Dahlbom, 1843: 147, junior homonym of *Harpactes* Swainson, 1833 (Aves), and of *Harpactes* Templeton, 1834 (Arachnida). Unjustified emendation of *Harpactus* Shuckard, 1837.

Dienoplus Fox, 1894: 548. Type species: *Dienoplus pictifrons* Fox, 1894, by monotypy.



Figs 1–2. Females habitus of *Harpactus* spp. (all from the Nakhchivan Autonomous Republic, Azerbaijan). 1 – *Harpactus vedicus* Nesterov, 1994; 2 – *H. elegans* (Lepeletier de Saint Fargeau, 1832).

Harpactus elegans (Lepeletier de Saint Fargeau, 1832)


Figs 2, 6–8

Arpactus elegans Lepeletier de Saint Fargeau, 1832: 79, ♂ (syntypes: ♂♂, France, Lyon area [depository unknown]).

Harpactus elegans: Nemkov, 1997: 19; Mokrousov *et al.*, 2019: 12.

MATERIAL EXAMINED. **Azerbaijan:** *Nakhchivan Autonomous Republic:* Babek, Gahab, 39°15'N 45°31'E, 1045 m, 12.VI 2019, 1 ♀; Babek, Payiz, 39°26'N 45°22'E, 1225 m, 11.VI 2019, 1 ♂; Shakhbuz, Zarnatun, 39°31'N 45°46'E, 1550 m, 14, 18.VI 2019, 2 ♂; Julfa, Gazanchi, 39°13'N 45°41'E, 1300 m, 15.VI 2019, 1 ♀, 6 ♂; Sharur, Akhura, 39°33'N 45°13'E, 1640 m, 13.VI 2019, 3 ♂; Shakhbuz, Kulus, 39°21'N 45°37'E, 1395 m, 19.VI 2020, 1 ♀.

DISTRIBUTION. North Africa (Algeria), Europe, Russia (European part, North Caucasus, Eastern Siberia), Armenia, Azerbaijan, Turkey, Kazakhstan.



Figs 3–8. *Harpactus* spp. (all from the Nakhchivan Autonomous Republic, Azerbaijan). 3–5 – *H. vedicus* Nesterov, 1994: 3 – female; 4, 5 – male; 6–8 – *H. elegans* (Lepeletier de Saint Fargeau, 1832): 6 – female; 7, 8 – male.

***Harpactus vedicus* Nesterov, 1994, stat. resurr.**

Figs 1, 3–5

Harpactus vedicus Nesterov, 1994: 35, ♀, ♂ (holotype: ♀, Armenia, Khozrov Nature Reserve, Vedic section, 1300 m, 6.VI.1982, M. Nesterov leg. [I.I. Schmalhausen Institute of Zoology of National Academy of Sciences of Ukraine, Kiev]), not examined. Synonymized with *Harpactus elegans* (Lepeletier de Saint Fargeau, 1832) by Nemkov, 1997: 19.

Harpactus transcaucasicus Nemkov, 1994: 69, ♀, ♂ (holotype, ♀: Azerbaijan, Nakhichevan ASSR, 35 km north of Nakhichevan, 20.VI.1985, V. Tobias leg. [ZISP]), examined. **Syn. n.**; Nemkov, 1996: 1208; Yildirim *et al.*, 2016: 16; Mokrousov *et al.*, 2019: 12; Kaplan & Yildirim 2021: 35; 2023: 1681.

MATERIAL EXAMINED. *Type material:* **Azerbaijan:** *Nakhchivan Autonomous Republic:* 35 km north of Nakhichevan, 20.VI 1985, 2 ♀, 12 ♂, V. Tobias (holotype and paratypes of *Harpactus transcaucasicus* Nemkov, 1994). *Other material:* **Azerbaijan,** *Nakhchivan Autonomous Republic:* Babek, 3 km NE Sirab, 39°18'N 45°32'E, 1250 m, 12.VI 2019, 1 ♀; Babek, Goynuk, 39°18'N 45°40'E, 1680 m, 12.VI 2019, 13 ♀, 15 ♂; Shakhbuz, Zarnatun, 39°31'N 45°46'E, 1550 m, 14, 18.VI 2019, 4 ♀, 33 ♂; Julfa, Gazanchi, 39°13'N 45°41'E, 1300 m, 15.VI 2019, 2 ♂; Julfa, Milakh, 39°15'N 45°43'E, 1430 m, 15.VI 2019, 1 ♀, 1 ♂; Julfa, Teyvaz, 39°15'N 45°46'E, 1645 m, 15.VI 2019, 2 ♀, 1 ♂; Ordubad, Aghdara, 39°06'N 45°54'E, 2000 m, 17.VI 2019, 1 ♂; Shakhbuz, Gomur, 39°27'N 45°44'E, 1790 m, 18.VI 2019, 1 ♀; Shakhbuz, Kechili, 39°22'N 45°43'E, 1800 m, 19.VI 2019, 1 ♀; Shakhbuz, Kulus, 39°21'N 45°37'E, 1395 m, 14, 19, 26.VI 2020, 6 ♀, 4 ♂; 27.VI 2021, 3 ♀; 4.VII 2021, 1 ♀, 1 ♂; Shakhbuz, Kechili, 39°22'N 45°43'E, 1800 m, 12.VII 2022, 6 ♂.

NOTE. Paratypes of *Harpactus vedicus* Nesterov, 1994 – 9 ♀, 90 ♂, Armenia, Khozrov Nature Reserve and 1 ♀, 1 ♂, Azerbaijan, Nakhchivan Autonomous Republic, Bilav could not be found in the ZISP collection, although according to Nesterov (1994: 35) some of them should be kept there.

DISTRIBUTION. Armenia, Azerbaijan, Turkey.

DISCUSSION

Harpactus vedicus Nesterov, 1994 was synonymized with *H. elegans* (Lepeletier de Saint Fargeau, 1832) by Nemkov (1997: 19), but the original description allowed conclusions to be drawn about the fallacy of this synonymy. P. Nemkov noted that *H. vedicus* differed from *H. elegans* only in the colouration of the body, and he considered it to be a colour form of the latter. We do not agree with this opinion, as the original description includes a comparison with *H. elegans*, which differs not only in colouration but also in the punctuation of the metasomal terga. Furthermore, the colouration characters of *H. vedicus* given by Nesterov in the original description are quite stable and clearly different from *H. elegans*: female face yellow with dark spot (dark medially in *H. elegans*, Fig. 3 vs. Fig. 6); yellow pattern on mesosoma more developed (Fig. 1 vs. Fig. 2); tergum I with large lateral spot (no spots or small unclear lateral spot on tergum I in *H. elegans*; Fig. 1 vs. Fig. 2); metasomal sterna III–VII of male dark (only sterna VI–VII dark in *H. elegans*; Fig. 5 vs. Fig. 8). Only material collected in the Arax valley (Nakhchivan Autonomous Republic, Azerbaijan) has been used for comparison purposes.

At the same time, *Harpactus vedicus* Nesterov and *H. transcaucasicus* Nemkov have the same colour and morphology. These facts, as well as the distribution (the Arax valley), lead to the conclusion that these species are conspecific. In addition, *H. elegans* is a relatively rare species in the Arax valley, while *H. vedicus* Nesterov is very common, as indicated by the number of type series, and this correlates well with the known abundance of *H. transcaucasicus*.

Both species (*Harpactus vedicus* Nesterov and *H. transcaucasicus* Nemkov) were described in the same year (1994), but the name *H. vedicus* Nesterov has priority because it was published earlier (February, 15) than *H. transcaucasicus* Nemkov (October, 14) (Article 23.1 – ICZN, 1999).

ACKNOWLEDGEMENTS

We are grateful to K. Fadeev (ZISP) for assistance in working with the ZISP collection, to A. Przhiboro (ZISP) for information on the date of publication of Nesterov's names, and to A. Lelej (FSCV) for valuable comments that helped improve the quality of this paper. The

research was carried out within the state assignment of Ministry of Science and Higher Education of the Russian Federation (theme No. 124012400285-7).

REFERENCES

- Bohart, R.M. & Menke, A.S. 1976. *Sphecid Wasps of the World. A generic revision*. University of California Press, Berkeley, Los Angeles and London. ix + 695 pp.
- Dahlbom, A.G. 1843. *Hymenoptera Europaea praecipue borealia; formis typicis nonnullis Specierum Generumve Exoticorum aut Extraneorum propter nexum systematicum associatis; per Familias, Genera, Species et Varietates disposita atque descripta. Tomus: Spheg in sensu Linneano*. Officina Lundbergiana, Lund. Fasc. 1: 1–172.
- Fox, W.J. 1894. The North American Larridae. *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1893: 467–551.
- International Commission on Zoological Nomenclature (1999) *International Code of Zoological Nomenclature. Fourth Edition adopted by the International Union of Biological Sciences*. The International Trust for Zoological Nomenclature, London. XXIX + 306 pp.
- Jurine, L. 1807. *Nouvelle méthode de classer les Hyménoptères et les Diptères. Hyménoptères. Tome premier*. J.J. Paschoud, Genève, 1–319 + [1] pp., 1–14 pls.
- Kaplan, E. & Yildirim, Y. 2021. An updated checklist of Turkish crabronid wasps (Hymenoptera: Crabronidae) with new and additional records. *Journal of Insect Biodiversity*, 21: 18–109. DOI: 10.12976/jib/2021.21.2.1
- Kaplan, E. & Yildirim, Y. 2023. A review of Sphecoformes (Hymenoptera) of Turkey, with history of their research. *Munis Entomology & Zoology*, 18: 1675–1710.
- Lepeletier de Saint Fargeau, A.L.M. 1832. Mémoire sur le G. *Gorytes* Latr. *Arpactus*. Jur. *Annales de la Société Entomologique de France*, 1: 52–79.
- Mokrousov, M.V., Proshchalykin, M.Yu., Aliyev, Kh.A. & Maharramov, M.M. 2019. To the knowledge of digger wasps (Hymenoptera: Crabronidae) of Nakhchivan Autonomous Republic of Azerbaijan. *Far Eastern Entomologist*, 394: 1–24. DOI: 10.25221/fee.394.1
- Nemkov, P.G. 1994 (14.X.1994). New species of digger wasps of the genus *Harpactus* (Hymenoptera, Sphecidae) from the central Palaearctic. *Zoologicheskii Zhurnal*, 73(11): 64–71. [In Russian]
- Nemkov, P.G. 1996. Digger wasps of the tribe Gorytini (Hymenoptera, Sphecidae) in the fauna of Russia and neighbouring countries. Genus *Harpactus* Shuckard. *Zoologicheskii Zhurnal*, 75(8): 1204–1213. [In Russian]
- Nemkov, P.G. 1997. To synonymies of Palaearctic digger wasps of the tribe Gorytini (Hymenoptera, Sphecidae). Part II. *Far Eastern Entomologist*, 47: 19.
- Nemkov, P.G. & Lelej, A.S. 2013. A cladistic analysis and classification of the subfamily Bembicinae (Hymenoptera: Crabronidae), with a key to the genera. *Zootaxa*, 3652(2): 201–231. DOI: 10.11646/zootaxa.3652.2.1
- Nesterov, M.A. 1994 (1993) (15.II.1994). Two new wasp species of the genus *Harpactus* (Hymenoptera, Sphecidae) from Armenia. *Journal of the Ukrainian Entomological Society*, 1(2): 35–40. [In Russian]
- Pulawski, W. 1985. *Harpactus* Shuckard, 1837, the valid name for the genus currently called *Dienoplus* Fox, 1894 (Hymenoptera: Sphecidae). *Systematic Entomology*, 10, 59–63.
- Pulawski, W. 2024. *Catalog of Sphecidae sensu lato*. Available from: <https://www.calacademy.org/scientists/projects/catalog-of-sphecidae> (Accessed 20 March 2024)

- Shuckard, W.E. 1837. *Essay on the indigenous fossorial Hymenoptera; comprising a description of all the British species of burrowing sand wasps contained in the metropolitan collections; with their habits as far as they have been observed*. Richter and Co., London. XII pp., one pl., 252 + [2] pp., pls. 1–4,
- Yildirim, E., Ljubomirov, T. Özbek, H. & Yüksel M. 2016. New data on Spheciformes fauna (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) of Turkey. *Journal of Insect Biodiversity*, 4: 1–51. DOI: 10.12976/jib/2016.4.3

© **Far Eastern entomologist (Far East. entomol.)** Journal published since October 1994.

Editor-in-Chief: S.Yu. Storozhenko

Editorial Board: A.S. Lelej, S.A. Belokobylskij, M.G. Ponomarenko, V.A. Mutin, E.A. Beljaev, E.A. Makarchenko, A.V. Gorochoy, T.M. Tiunova, M.Yu. Proshchalykin, S.A. Shabalin, V.M. Loktionov

Address: Federal Scientific Center of the East Asia Terrestrial Biodiversity (former Institute of Biology and Soil Science), Far East Branch of the Russian Academy of Sciences, 690022, Vladivostok-22, Russia.

E-mail: storozhenko@biosoil.ru

web-site: <http://www.biosoil.ru/fee>