

A NEW SPECIES OF *MANINEURA* BOUČEK, 1979 (CHALCIDOIDEA: PTEROMALIDAE) FROM THE PALAEARCTIC REGION

E. V. Tselikh^{1,*}, J. Lee², D-S. Ku³

1) Zoological Institute, Russian Academy of Sciences, Universitetskaya nab. 1, St. Petersburg 199034, Russia. *Corresponding author, E-mail: tselikhk@gmail.com

2) Department of Plant Medicine, Gyeongsang National University, Jinju 52828, Republic of Korea E-mail: wogus0913@naver.com

3) The Science Museum of Natural Enemies, Geochang 50147, South Korea. E-mail: bracon2700@hanmail.net

Summary. The rare genus *Manineura* Bouček, 1979 is recorded for the first time from the Palaearctic region. A new species *Manineura deokseoi* sp. n. is described and illustrated from South Korea. An updated diagnosis of the genus *Manineura* is given.

Key words: Chalcidoidea, Pteromalidae, Metasteninae, taxonomy, new species, new records, Palaearctic region.

Е. В. Целих, Дж. Ли, Д-С. Ку. Новый вид *Manineura* Bouček, 1979 (Chalcidoidea: Pteromalidae) из Палеарктики // Дальневосточный энтомолог. 2026. N 543. С. 10-16.

Резюме. Впервые в Палеарктическом регионе обнаружен род *Manineura* Bouček, 1979. Описан и проиллюстрирован новый вид *Manineura deokseoi* sp. n. из Южной Кореи. Приведен обновленный диагноз для рода *Manineura*.

INTRODUCTION

In the last decade, a lot of papers have been devoted to the family Pteromalidae (Hymenoptera: Chalcidoidea) of the Eastern Palaearctic region, including descriptions of new taxa from Korea and adjacent regions (Tselikh, 2019; Tselikh *et al.*, 2022, 2023, 2024, 2025a, 2025b, 2025c; Li *et al.*, 2025).

The pteromalid genus *Manineura* Bouček, 1979 belongs to the subfamily Metasteninae (Burks *et al.*, 2022). Up to now, it consists of only one species, *Manineura pentatomivora* (Mani, 1939), known from the Oriental region (UCD Community, 2026). Although the biology of this genus has not been studied in detail, it is known that *M. pentatomivora* is the primary parasitoid of *Urostylis punctigera* Westwood, 1837 (Hemiptera: Pentatomidae).

During our study of Pteromalidae in Asia (Tselikh *et al.*, 2022, 2023, 2024, 2025a, 2025b, 2025c), several specimens of a new *Manineura* species were collected in forested areas of South Korea. These specimens represent the first recorded occurrence of the genus in the Palaearctic region. Here, we describe this new species, *Manineura deokseoi* sp. n., and provide a comparative diagnosis of this species with described species of *Manineura*.

MATERIAL AND METHODS

Morphological terminology, including sculpture and wing venation, follows Bouček and Rasplus (1991), Gibson (1997), and Burks *et al.* (2022). The following abbreviations are used: POL – posterior ocellar line, the minimum distance between the posterior ocelli; OOL – ocello-ocular line, the minimum distance between a posterior ocellus and compound eye; C1–C4 – clavomeres; M – marginal vein; S – stigmal vein; PM – postmarginal vein; F1–F6 – funicular segments. The scape is measured without the radicle; the pedicel is measured in lateral view. The distance between the lower margin of the clypeus and the toruli is measured from the lower margins of the toruli. Eye height is measured as the maximum diameter, eye length as the minimum diameter. Mesosoma and metasoma are measured in lateral view, the latter including the ovipositor sheaths.

Specimens examined in this paper are deposited in the Hymenoptera collections of the National Institute of Biological Resources, Incheon, Republic of Korea (NIBR), the Zoological Institute of the Russian Academy of Sciences, St. Petersburg, Russia (ZISP) and the Natural History Museum, London, England (NHMUK).

TAXONOMY

Family Pteromalidae

Subfamily Metasteninae

Genus *Manineura* Bouček, 1979

Manineura Bouček, 1979: 471–472.


Type species: *Pachyneuron pentatomivora* Mani, 1939, by original designation.

DIAGNOSIS. Head in frontal view high as broad, lower face convex and long (Figs 2, 10). In dorsal view strongly convex, without occipital carina (Figs 6, 14). Lower margin of clypeus slightly produced in middle (Fig. 11) or rounded (Fig. 4). Antennal formula 11264; anelli small, microsetose on C3–C4 (Figs 5, 13). Antennal toruli situated above level of lower edges of eyes (antenna high above center of face); antennal scrobes not deep (Figs 2, 10). Mandibles large and falcate (Fig. 2); mouth margin in anterior view broad-truncate and distinctly wavy.


Mesosoma short, strongly arched (Figs 1, 9). Pronotum very short and hardly visible; collar margin without carina (Figs 6, 7, 14). Notauli incomplete (Figs 7, 8, 14). Axilla not strongly advanced; axillula not enlarged. Scutellum arched, without conspicuous sublateral grooves, not distinct frenal area and without frenal groove (Fig. 8). Propodeum transverse, short and alutaceous or smooth, without costula, median carina and plicae, nucha short, propodeal spiracles near to front margin of sclerite (Fig. 8). Fore wing hyaline; marginal vein abruptly thickened at parastigmal break relative to submarginal vein thickness (Figs 4, 6, 15), M wide and short, stigma large (Figs 3, 12). Hind coxa dorsally bare; hind tibia with one spur.

Metasoma shorter than combined length of mesosoma and head (Figs 1, 9). Petiole distinct, anteroventrally either braced by a flange extending from the 1st gastral sternum that reaches anteriorly under the petiolar attachment. Cerci with setae subequal in length. Ovipositor not strongly protruding.

DISTRIBUTION. Palearctic (new record) and Oriental regions (UCD Community, 2026).



Figs 1–8. *Manineura mavca* **sp. n.**, holotype, female: 1 – body, lateral view; 2 – head, frontal view; 3 – fore wing; 4 – clypeus; 5 – antenna; 6 – head and pronotum, dorsal view; 7 – body, dorsal view; 8 – mesosoma, dorsal view.



Figs 9–15. *Manineura pentatomivora* (Mani, 1939), paratype, female: 9 – body, lateral view; 10 – head, frontal view; 11 – clypeus; 12 – fore wing; 13 – antenna; 14 – body, dorsal view; 15 – head and mesosoma, lateral view.

ACKNOWLEDGEMENTS

This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Climate, Energy and Environment of the Republic of Korea (NIBR202304203, NIBR202402202, NIBR202502202). And it was partially funded by Russian State Research (project No. 125012901042-9).

The authors are very thankful to Natalie Dale-Skey, she provided comprehensive assistance with our work at the Natural History Museum. We are grateful to all referees for their valuable comments.

REFERENCES

- Bouček, Z. & Rasplus, J.-Y. 1991. *Illustrated key to West-Palaeartic genera of Pteromalidae (Hymenoptera: Chalcidoidea)*. Institut National de la Recherche Agronomique, Paris. 140 pp.
- Bouček, Z. 1979. *Oricoruna* and *Manineura*, new pteromalid genera (Hymenoptera) from the oriental region. *Oriental Insects*, 12(4): 469–472.
- Burks, R., Mitroiu, M.-D., Fusu, L., Heraty, J.M., Janšta, P., Heydon, S., Dale-Skey Papilloud, N., Peters, R.S., Tselikh, E.V., Woolley, J.B., Noort, S., Baur, H., Cruaud, A., Darling, Ch., Haas, M., Hanson, P., Krogmann, L., Rasplus, J.-Y. 2022. From hell's heart I stab at thee! A determined approach towards a monophyletic Pteromalidae and reclassification of Chalcidoidea (Hymenoptera). *Journal of Hymenoptera Research*, 94: 13–88. DOI: 10.3897/jhr.94.94263
- Gibson, G. 1997. Morphology and Terminology. P. 16–44. In: Gibson, G.A.P, Huber, J.T., Woolley, J.B. (Eds.). *Annotated Keys to the Genera of Nearctic Chalcidoidea (Hymenoptera)*. NRC Research Press, Ottawa.
- Li, X., Yin, H., Li, Q. & Tselikh, E.V. 2025. New data on *Halticoptera* Spinola, 1811 and *Sphegigaster* Spinola, 1811 (Chalcidoidea: Pteromalidae) from China. *Far Eastern Entomologist*, 527: 11–16. DOI: 10.25221/fee.527.2
- Mani, M.S. 1939. Descriptions of new and records of some known chalcidoid and other hymenopterous parasites from India. *Indian Journal of Entomology*, 1: 69–99.
- Tselikh, E.V. 2019. New records of the chalcid wasps of the family Pteromalidae (Hymenoptera: Chalcidoidea) from Russia. *Far Eastern Entomologist*, 385: 12–24. DOI: 10.25221/fee.385.2
- Tselikh, E.V., Lee, J. & Ku, D-S. 2022. New records of the chalcid wasps of the family Pteromalidae (Hymenoptera: Chalcidoidea) from South Korea. *Far Eastern Entomologist*, 462: 8–19. DOI: 10.25221/fee.462.2
- Tselikh, E.V., Lee, J. & Ku, D.-S. 2023. New data on the chalcid wasps of the family Pteromalidae (Hymenoptera: Chalcidoidea) from South Korea. *Far Eastern Entomologist*, 482: 10–21. DOI: 10.25221/fee.482.2
- Tselikh, E.V., Lee, J. & Ku, D-S. 2024. Review of the South Korean species of *Plutothrix* Förster, 1856 (Hymenoptera: Pteromalidae), with description of three new species. *Far Eastern Entomologist*, 508: 1–18. DOI: 10.25221/fee.508.1
- Tselikh, E.V., Rasplus, J.-Y., Lee, J. & Ku, D-S. 2025a. *Vedma circe* (Hymenoptera: Chalcidoidea: Pteromalidae), a new genus and species from South Korea. *Zoosystematica Rossica*, 34(2): 285–288. DOI: 10.31610/zsr/2025.34.2.285
- Tselikh, E.V., Rasplus, J.-Y., Lee, J. & Ku, D-S. 2025b. *Krasava*, a new genus of Pteromalidae (Hymenoptera, Chalcidoidea) from the Palaeartic region. *Zootaxa*, 5715(1): 489–493. DOI: 10.11646/zootaxa.5715.1.42

Tselikh, E.V., Rasplus, J.-Y., Lee, J. & Ku, D-S. 2025c. A new species of *Uniclypea* Bouček, 1976 (Chalcidoidea, Pteromalidae) from the Palearctic region. *Far Eastern Entomologist*, 523: 1–4. DOI: 10.25221/fee.523.1

UCD Community. 2026. Universal Chalcidoidea Database (UCD) curated in TaxonWorks. 1143. Available at: <https://sfg.taxonworks.org/api/v1/> (Accessed on 10 January 2026)