

Chromosomal forms of *Microtus maximowiczii* (Schrenck, 1859) (Rodentia, Cricetidae): variability in 2n and NF in different geographic regions

Irina V. Kartavtseva*, Irina N. Sheremetyeva, Kira V. Korobitsina, Galina A. Nemkova, Elena V. Konovalova, Vladimir V. Korablev & Leonid L. Voyta

ABSTRACT. Analysis of the chromosomal characteristics of the Maximowicz's vole *Microtus maximowiczii* (242 specimens) from Transbaikalia, Russian Far East, and Mongolia, including new data from 12 local populations, supplemented information about chromosomal polymorphism of the species ($2n=36-44$; $NF=52-62$). We describe chromosomal data for the populations which have not been investigated previously, reveal a fixation of the chromosomal reconstructions in the various geographical regions and describe for the first time the population with stabilized karyotype ($2n=42$) in the Chita Region. The revision of the chromosomal characteristics has been done, that resulted in suggestion to consider five forms in the species composition, three of which (C, D and I) are established herein. Perhaps *M. m. gromovi* exemplified by the chromosome form I should be considered as an independent species. The chromosomal forms differ not only in 2n and NF, but also in number of the biarmed and acrocentric chromosomes.

KEY WORDS: chromosome, karyotype, polymorphism, *Microtus maximowiczii*.

Irina V. Kartavtseva [Irina-kar52@rambler.ru], Irina N. Sheremetyeva, Kira V. Korobitsina, Galina A. Nemkova, Elena V. Konovalova, and Vladimir V. Korablev, Institute of Biology and Soil Science of the Far East Branch of the Russian Academy of Sciences, Vladivostok 690022, Russia; Leonid L. Voyta, Institute of Natural Resources, Siberian Branch of Russian Academy of Sciences, Chita 672014, Russia.

Хромосомные формы *Microtus maximowiczii* (Schrenck, 1859) (Rodentia, Cricetidae): изменчивость 2n и NF в различных географических регионах

И.В. Картавецва, И.Н. Шереметьева, К.В. Коробицына, Г.А. Немкова,
Е.В. Коновалова, В.П. Кораблев, Л.Л. Войта

РЕЗЮМЕ. Анализ хромосомных характеристик 242 полевок Максимовича из Забайкалья, Дальнего Востока России и Монголии, включая собственные новые данные из 12 локальных популяций, позволил дополнить информацию о хромосомном полиморфизме вида ($2n=36-44$; $NF=52-62$), описать хромосомные данные для ранее не исследованных популяций, выявить фиксацию хромосомных перестроек в различных географических регионах, впервые описать популяцию со стабилизировавшимся кариотипом ($2n=42$) в Читинской области. В результате ревизии хромосомных характеристик выделено шесть форм, три из которых (С, D и I) установлены впервые. Подвид *M. m. gromovi*, представленный хромосомной формой I, возможно, следует рассматривать как самостоятельный вид. Показано, что хромосомными характеристиками для каждой формы являются не только 2n и NF, но и числа двуплечих и акроцентрических хромосом.

КЛЮЧЕВЫЕ СЛОВА: хромосома, кариотип, полиморфизм, *Microtus maximowiczii*.

Introduction

The Maximowicz's vole *Microtus maximowiczii* (Schrenck, 1859) is a widely distributed species inhabiting humid biotopes of the forest zone in East Asia (Russia, Mongolia, and North-East China). For the Russian territory two subspecies have been described: *M.*

m. unguensis Kastschenko, 1912 in Transbaikalia and *M. m. maximowiczii* (Schrenck, 1859) in Amur Region. The polymorphism in number and morphology of chromosomes ($2n=36-44$; $NF=52-62$) for this species has been reported before for 134 animals from 19 local populations (Kovalskaya, 1977; Kovalskaya *et al.*, 1980; Golenishchev & Radjabli, 1981; Meyer *et al.*, 1996). Three chromosomal forms (A, B, and V) based on 2n and NF was shown for this area (Kovalskaya *et al.*,

* Corresponding author




Figure 1. Geographic position of *Microtus maximowiczii* (Schrenck, 1859) samples (numbers correspond to those in tab. 1).

1980). The chromosomal form “A” with high chromosomal numbers ($2n=42-44$; $NF=56-62$) includes individuals from the Northern and Central Transbaikalia and Amur area. Voles of the chromosomal form “B” with $2n=38-42$ are noted in the lower reaches of Selenga River ($2n=39-41$; $NF=54-57$) and in the middle part of the Onon River in the Amur River basin ($2n=39-42$; $NF=54-58$). The voles of the chromosomal form “V” with the least numbers of chromosomes ($2n=36-38$; $NF=52-55$) are found out on slopes of Hentey Mountains in Mongolia (Kovalskaya *et al.*, 1980). The G-banding of several animals of two chromosomal forms makes possible to show, that chromosomal polymorphism of *M. maximowiczii* is connected with Robertsonian reorganizations, inversions and also tandem fusion of middle metacentrics with formation of large metacentrics (Meyer *et al.*, 1996). According to data of the chromosomal reorganizations proved by using of methods of differential staining of chromosomes, the number of chromosomal rearrangements in various geographic populations should be reconsidered. Therefore nowadays the chromosomal variability of *M. maximowiczii* is mainly still unknown. The purpose of the present work is to investigate variability of chromosomal characteristics: $2n$, NF , number of biarmed (meta-, submeta-, and subtelocentrics) and acrocentric chro-

mosomes in karyotype of Maximowicz’s vole throughout its area, using both original and published data.

Material and methods

The karyotypes of 242 animals from 40 local populations based on original and published data were investigated (Fig. 1; Tab. 1). We present here new chromosomal data of 94 voles from 12 localities: 76 Maximowicz’s voles from ten new localities (16, 21, 22, 26–28, 30, 31–32, and 33) and 18 voles from two localities (5 and No. 12), from which chromosomal data were known earlier, have been trapped in 2003–2008. The coordinates of localities were provided only for our data (Tab. 1). Chromosomes were obtained from bone marrow cells. After colchicine (0.04%) treatment and hypotonic treatment with KCl-solution (0.56%), the cells were fixed with acetic-methanol (1:3), then air or flame-dried and stained with 4% Giemsa. The preparations were processed in the field with exception of a few animals. The diploid number ($2n$) and total number of chromosomal arms (fundamental number, NF) were determined after examination about 20 metaphases per animal. The chromosomal forms are given in accordance to the characteristics proposed by Kovalskaya (1977).

Table. Chromosomal data of *Microtus maximowiczii* (Schrenk, 1859) from different regions

Locality and specimen numbers (##)		2n	Chromosomes		NF	Reference
			Biarmed M+SM+ST	A		
<i>Microtus maximowiczii unguurensis</i> Kastschenko, 1912						
Form B						
1	Buryatia, Kabansk District, Baikalo-Kudara (n=5)					1
	#48	38	16	22	54	
	#47, #503	40	14	26	54	
	#385	40	15	25	55	
	#226, #227	41	15	26	56	
2	Buryatia, Selenga River delta, Sotnikovskii Island (n=3)					
	#78	40	15	25	55	
	#186	41	?	?	55	
	#182	41	16	25	57	
3	Buryatia, Selenga River delta, Kolesovskogo Island (n=7)					
	#159, #158	38	17	21	56	
	#81, #79, #80	39	16	23	55	
	#196	40	15	25	55	
	#187	40	16	24	56	
4	Buryatia, Kabansk District, Istomino (n=7)					
	#362, #596, #615	39	16	23	55	
	#361	41	14	27	55	
	#572, #571, #529	40	16	24	56	
5	Ibidem (2006), N 52008' 26.00", E 106017' 57.85" (n=5)					our data
	#2023	38	19	19	57	
	#2016	39	17	22	56	
	#2031	40	16	24	56	
	#2017, #2030	40	14	26	54	
	#2024	42	14	28	56	
Total		38–42	14–19	19–28	54–57	
Form A						
6	Buryatia, Bauntovsk District, Bagdarino (n=6)					1
	#503	40	14	26	54	
	#569	42	14	28	56	
	#625	41	16	26	58	
	#606	41	17	25	58	
	#598	41	15	26	57	
	#595	43	16	27	59	
7	Buryatia, Bauntovsk District, Romanovka (n=2)					2
	no specimen number	42	16	26	58	
	no specimen number	43	17	26	60	
8	Ibidem (n=2)					3
	no specimen number	42	14	28	56	
	no specimen number	43	15	28	58	
9	Ibidem (n=2)					1
	#407, #433	42	16	26	58	
10	Ibidem (n=1)					4
	no specimen number	42	16	26	58	
11	Ibidem (n=4)					1
	#566	44	18	26	62	
	#415	42	16	26	58	
	#504	42	15	27	57	
	#221	42	14	28	56	
	4 hybrids (#433 x #566: 2n=42 x 2n=44)	43	17	26	60	

Table (continued).

Locality and specimen numbers (##)		2n	Chromosomes		NF	Reference
			Biarmed M+SM+ST	A		
<i>Microtus maximowiczii unguensis</i> Kastschenko, 1912 Form B						
12	Ibidem (n=13) #2350 #2375, #2335, #2336, #2349, #2353, #2373 #2373 #2374, #2333 #2334 #2345 #2352	41 42 43 43 43 40 44	16 15 17 16 13 13 18	25 27 26 27 30 27 26	57 57 60 59 56 53 62	our data
13	Buryatia, 50 km from Romanovka, Endodinka River (n=2) no specimen number no specimen number	42 43	14 15	28 28	56 58	1
14	Transbaikalia, Ivan Lake (n=2) no specimen number no specimen number	43 44	17 16	26 28	60 60	2
15	Transbaikalia, Ivan Lake (n=3) no specimen number no specimen number	42 44	14 16	28 28	56 60	3
16	Transbaikalia, Malyi Undugun Lake (n=17) #1754 #1737, #1744, #1745, #1750, #1752, #1739 #1749, #1759 #1738, #1740, #1741, #1743, #1747, #1751 #1742 #1748	41 42 42 43 43 43	15 15 13 15 16 14	26 27 29 28 27 29	56 57 55 58 59 57	our data*
17	Transbaikalia, Beklemishevo (n=5) no specimen numbers	42–43	□	□	55–59	5
18	Transbaikalia, near Chita city (n=3) no specimen numbers	41–42	15–16	26	52–56	2
19	Transbaikalia, Makkaveevo (n=21) no specimen numbers one animal from this sample without number	42–43 42	□ 16	□ 26	56–59 58	5
20	Transbaikalia, Akshinsk District, Ureisk (n=15) no specimen numbers one animal from this sample without number	39–42 42	13–18 17	21–29 24	54–58 58	
	Total	39–44	13–18	21–30	53–60	
Form V						
21	Transbaikalia, Sokhondinskii Reserve, Bukukun River, N 49027' 40", E 111007' 11.36" (n=7)* #1668, #1678, #1682 #1667, #1688, #1694 #1669	38 39 40	16 16 16	22 23 24	54 55 56	our data*
22	Ibidem, Enda River, N 49027' 36.74" E 1110021' 22.10" (n=12) #1671 #1672 #1665, #1666, #1685, #1662, #1676, #1697, #1692 #1663, #1664 #1696	36 37 38 39 40 40	16 15 16 17 16 □	20 22 22 22 24 □	52 52 54 56 56 □	
23	Mongolia, Hentei Aimak, Bat-Shiret, Onon River (n=13) no specimen numbers	36–38	14–18	18–24	52–55	5
24	Mongolia, Central Aimak Dzynbyrx and Borxon rivers (n=21) no specimen numbers one animal from this sample without number	36–37 36	15–18 17	18–22 19	52–54 53	
	Total	36–40	15–18	18–24	52–56	

Table (continued).

Locality and specimen numbers (##)		2n	Chromosomes		NF	Reference
			Biarmed M+SM+ST	A		
<i>Microtus maximowiczii unguurensis</i> Kastschenko, 1912						
Form D						
25	Mongolia, East Aimak, Halhin-Gol River (n=3)					5
	no specimen numbers	42–43	14–15	28	56, 58	
26	Transbaikalia, Krasnokamenskii Reserve, near Krasnokamensk (n=2)					our data
	#2120	40	14	26	54	
	#2121	43	15	28	58	
27	Transbaikalia, Kovyli (n=1)					our data**
	#2343	42	14	28	56	
28	Transbaikalia, Baleiskii reserve, Podoinicino, N 51040' 26.77", E 116047' 28.86" (n=8)					
	#2326–2328, #2330–2332, #2347–2348	42	14	28	56	
	Total	40–43	14–15	26–28	54–58	
<i>Microtus maximowiczii maximowiczii</i> (Schrenck, 1859)						
Form C						
29	Amur Region, Klimoutci (n=1)					6
	no specimen number	42	20	22	62	
30	Amur Region, near Blagoveshchensk, N 50015' 22.13", E 127034' 41.58V (n=1)					our data
	# 1659	42	16	26	58	
31	Amur Region, Norskii reserve (2001), N 52029.019', E130000.948' (n=18)					our data**
	#1518–1522, #1526, #1528, #1531–1533, #1537, #1539	40	18	22	58	
	#1527, #1529–1530, #1536, #1538	41	19	22	60	
32	Ibidem (2006), N 52029.019', E 130000.948' (n=8)					our data
	#2060, #2064–2066	40	18	22	58	
	#2059, #2061–2063	41	19	22	60	
33	Amur Region, Arhara, N 49006' 11.46", E 130006' 11.41" (n=1)					our data
	#2058	41	19	22	60	
34	Evreiskii Avtonomnyi Region, near Birabidjan (n=1)					7
	no specimen number	40	18	22	58	
35	Evreiskii AR, Aur (n=1)					
	no specimen number	41	18	22	58	
36	Evreiskii AR, Leninskoe (n=4)					2
	no specimen numbers	40	18–19	22	58–60	
	no specimen number	41	19	22	60	
37	Khabarovsk Territory, Utinaya station (n=5)					2
	no specimen numbers	39–41	16–18	22–23	55–59	
	one animal from this sample without number	41	18	23	59	
	Total	39–42	16–19	22–63	55–60	
<i>Microtus maximowiczii gromovi</i> Vorontsov, Boeskorov, Lyapunova & Revin, 1988						
Form I						
38	Yakutia, Bolshoe Toko Lake (n=1)					8
	# S-140238	44	16	28	60	
39	Khabarovsk Territory, Ayano-Maisk District, Ayan (2002; n=3)					9
	##1453–1455	44	16	28	60	
40	Ibidem (2008; n= 1)					our data
	#2383	44	16	28	60	
	Total	44	16	28	60	

* Published previously in Korobitsyna *et al.*, 2005.

** Published previously in Kartavtseva *et al.*, 2007.

References: 1) Kovalskaya, 1977; 2) Meyer *et al.*, 1996; 3) Golenishchev & Radjabli, 1981; 4) Meyer, 1978; 5) Kovalskaya *et al.*, 1980; 6) Meyer *et al.*, 1967; 7) Frisman *et al.*, in press; 8) Vorontsov *et al.*, 1988; 9) Sheremetyeva *et al.*, 2008.




Figure 2. Karyotypes of *Microtus maximowiczii unguensis* Kastschenko, 1912 from Istomino village, Buryatia. The united chromosomes are underlined.

A — #2016, $2n=39$; B — #2017, $2n=42$; C — #2024, $2n=42$; D — #2031, $2n=40$.

Results and discussion

New chromosomal data ($2n$ and NF) of 94 voles from 12 localities and literature data of three *M. maximowiczii* subspecies investigated from Transbaikalia and the Russian Far East is presented in the Tab. 1 and Fig. 1.

Microtus maximowiczii unguensis Kastschenko, 1912

Form B. Buryatia, Baikal Lake shore, near delta of Selenga River (localities 1–5). For animals from Istomino village (locality 5) we can see polymorphism connected with three variants of chromosomal rearrangement of the sixth pair of chromosomes: two metacentrics and four acrocentrics with two Robertsonian reorganizations and one fusion of metacentrics with formation of large metacentrics (Fig. 2).

Six animals from this sample have $2n=38–40$, 42 chromosomes with NF=54–56–57: two animals (#2017

and #2030) have the same chromosomal data ($2n=40$, NF=54) (Fig. 2A) and four — different: #2016 — $2n=39$, NF=56 (Fig. B); #2024 — $2n=42$, NF=56 (Fig. 2C); #2031 — $2n=40$, NF=56 (Fig. 2D) and #2023 — $2n=38$, NF=57.

Karyotypes of eight animals from this sample have been described before (Tab. 1, locality 4) with $2n=39–41$ and NF=55–57, but with different variants of the sets with three Robertsonian rearrangement and one pericentric inversion (Kovalskaya, 1977). We have confirmed the chromosome fusion with formation of three pairs of metacentric. But Kovalskaya (1977) presumed formation of all metacentrics by acrocentric chromosomes. The same chromosomal variant with $2n=40$, NF=54–56 has been found in Buryatia: Kabansk District, Baikalo-Kudara village, locality 1 and Bauntovsk District, Bagdarino village, locality 6 ($2n=40$, NF=54–59; Tab. 1). In other samples (localities 2 and 3) from delta of Selenga River we presume the same variants of rearrangements and stabilization of karyotype in re-

spect of two large metacentrics in Kolesovskogo Island (locality 3) (Kovalskaya, 1977).

Form A. Central Transbaikalia, Buryatia, near Romanovka village (locality 12). We investigated chromosomal sets of 13 animals with $2n=40-44$ and $NF=52-62$. From this locality, eight animals have been described before (7–11) with $2n=42-44$ and $NF=57-62$ (Kovalskaya, 1977; Meyer, 1978; Golenishchev & Radjabli, 1981; Meyer *et al.*, 1996). The karyotype of *M. maximowiczii* with $2n=44$ was described twice: from this place and from sample 14. We have found out one more animal with $2n=44$ in the sampling 12. The form A has $2n=42-44$ according to Kovalskaya (1977), but we have trapped animal also with $2n=40$.

Transbaikalia, Malyi Undugun Lake (locality 16). Chromosomal sets of 18 animals were: $2n=41-43$, $NF=57-59$ from which five animals have $2n=42$, $NF=57$ and six animals $2n=43$, $NF=58$. Preliminary chromosomal data with $2n=41-43$, $NF=55-59$ from this locality has been published earlier (Korobitsyna *et al.*, 2005). We assume a similarity of chromosomal characteristics of animals from two samples from Ivan Lake (localities 14–15) (Golenishchev & Radjabli, 1981; Meyer *et al.*, 1996) which is located close to the Malyi Undugun Lake.

The published data about chromosomal sets of animals from the localities in Chita Region (localities 17–20) allow presuming the same variants of chromosomal rearrangements like in sample 16. Our specimens can be referred to Form A.

Form V. Transbaikalia, Sokhondinskii Reserve (localities 21–22). Two local populations isolated from each other by a large forest and mountain range have been investigated near southern border of the Sokhondinskii Reserve. The first population is localized in the lower reach of the Bukukun River (locality 21). From seven individuals studied, three had $2n=38$, $NF=54$, three $2n=39$, $NF=55$, and one $2n=40$, $NF=56$. The total chromosomal data for this population is $2n=38-40$, $NF=54-56$.

The second population is localized in the lower reach of the Enda River (locality 22). The total chromosomal data in a small population, with the greater spectrum of chromosomal variability was found ($2n=36-40$, $NF=52, 54, 56$). Low value of chromosomal number ($2n=36$) in this population is similar to those found in the three localities from Mongolia ($2n=36-37$, $NF=52-54$, locality 23 and $2n=36-37$, $NF=52-54$, locality 24; Tab. 1). Our data coincides with chromosomal characters of the form V.

Form D. South-East Transbaikalia (localities 26–28). From this region we investigated chromosomal sets of grey voles from three local populations:

1) Chromosomal sets of two animals from locality near Krasnokamensk city (locality 26) were $2n=40$, $NF=54$ and $2n=43$, $NF=58$.

2) One animal, trapped near Kovyli village (locality 27), had $2n=42$, $NF=56$.

3) Near Podoinicino village (locality 28) all animals (not only from one family) had $2n=42$, $NF=56$: 6 SM-ST, 2 M (M1/M1), 6 middle M and 26 A, X — middle

A, Y — small A (Fig. 3A). This kind of karyotype was typical for many animals from Chita Region (localities 14–19) and Halhin-Gol in Mongolia (locality 25).

Hitherto it has not been described any population of *M. maximowiczii* with stable karyotype. This means that in the given population (locality 28) it occurred a stabilization of the karyotype described above. Such variant of karyotype is noted also in the population from South-East Mongolia, East Aimak, Halhin-Gol River (locality 25). We do not exclude an opportunities of karyotype stabilization in populations in future. Just this chromosomal variant in the Chita Region populations confirms that the similar chromosome pairs participate in the reconstruction in these regions. This set of the chromosomal rearrangements has not been found in the voles of form A (our data), therefore we propose for this the new Form D.

Microtus maximowiczii maximowiczii (Schrenck, 1859)

Form C. Amur area (localities 30–33). In the Amur area we investigated chromosomal sets of 28 animals from three local populations:

1) One animal of *M. maximowiczii* with $2n=42$, $NF=58$ has been trapped inside the *M. fortis* colony near Blagoveshchensk city (locality 30).

2) Twenty six animals from the Norskii Reserve, the left coast of the Nora River, had $2n=40-41$, $NF=58-60$. We trapped animals during two years, 2001 (locality 31) and 2006 (locality 32). This population is characterized by polymorphism: only by one rearrangement: fusion of two middle sized pairs of metacentrics up to formation large metacentric.

3) One animal from Arhara urban village (locality 33) had $2n=41$, $NF=60$. This variant of karyotype is similar with karyotypes of voles from the Norskii Reserve. The G-banding of chromosomes was applied for voles from the Norskii Reserve and Arhara urban village that permitted to determine the pairs involved in the rearrangements mentioned above (Fig. 3B). In this geographical region we deal with chromosomal polymorphism resulted from three reconstruction: fusion of the metacentric chromosomes with formation of large metacentric chromosome, fusion of the acrocentric chromosomes from two pairs with formation a submetacentric chromosome of the middle size, and pericentric inversion of the small acrocentric with formation of small subtolocentrics. Fixation of the chromosomal rearrangements in the vole populations of this region should be studied additionally. This combination of the chromosomal rearrangements has not been found in the voles of form A (our data), so they are attributed to the new Form C.

Microtus maximowiczii gromovi Vorontsov, Boeskov, Lyapunova & Revin, 1988

Form I. Khabarovsk Territory, Ayano-Maisk District (localities 39 and 40). The karyotype of *M. m. gromovi* has no any big metacentrics and differs from



Figure 3. A — Chromosomal set of *M. maximowiczii unguurensis* Kastschenko, 1912 (male #2327) from Podoinicino village, 2n=42. B — G-banding of chromosomes of *M. maximowiczii maximowiczii* (Schrenk, 1859) (female #2058) from Arhara village.

the simple chromosomal variant (2n=44, NF=60) by pericentric inversions on three St pairs of chromosomes including one large St pair (G banding was applied), which in *M. m. maximowiczii* and *M. m. unguurensis* we always meet in constant size and morphology, while in *M. m. gromovi* this pair is presented by a large acrocentric. The data on the morphological and chromosomal analyses enables to consider *M. m. gromovi* (form I) as an independent species (Kartavtseva *et al.*, 2008; Sheremetyeva *et al.*, 2008).

Conclusion

The published and presented here data confirms a high level of chromosomal polymorphism in *M. maximowiczii* (2n=36–44; NF=52–62). There are no less than five chromosomal forms but not three as it has been supposed before. These forms are characterized by various combinations of chromosomal rearrangements with different number of biarmed and acrocentric chromosomes. Characteristics for chromosomal forms are: Form A: 2n=39–44, NF=53–60 (D=13–18, A=21–30); Form B: 2n=38–42, NF=54–57 (D=14–19, A=19–28); Form C: 2n=39–42, NF=55–60 (D=16–19, A=22–26); Form V: 2n=36–40, NF=52–56 (D=15–18, A=18–24); Form D: 2n=40–43, NF=54–58 (D=14–15, A=26–28); Form I: 2n=44, NF=60 (D=16, A=28). It seems, in each geographic region peculiar chromosomal rearrangements and their combinations tend to be fixed. The using of methods of differential staining and ZooFISH of chromosomes will allow defining distinctly the characters of these rearrangements in the chromosomal forms. Clear chromosomal differentiation of *M. m. maximowiczii* from *M. m. unguurensis* is shown. We also confirm species status of subspecies *M. m. gromovi*.

ACKNOWLEDGEMENTS. We are very grateful to the colleagues from Khinganskii, Norskii, and Sokhondinskii

reserves for trapping of animals. The study was supported by the RFBR (## 06-04-48969 and 07-04-10175) and FEB RAS #06-III-A-06-473 grants.

References

- Frisman L.V., Korobitsyna K.V., Kartavtseva I.V., Sheremetyeva I.N. & Voyta L.L. 2009. Grey voles from the Russian Far East allozyme and karyological divergence // Russian Journal of Genetics. Vol.45. No.6. P.707–714.
- Golenishchev F.N. & Radjabli S.I. 1981. [New species of gray vole from shore of Evoron Lake] // Doklady Akademii Nauk SSSR. Vol.257. No.1. P.248–250 [in Russian].
- Kartavtseva I.V., Korobitsyna K.V., Sheremetyeva I.N., Nemkova G.A., Konovalova E.V., Korablev V.P., Frisman L.V., Kryukov A.P. & Voita L.L. 2008. Chromosomal polymorphism of *Microtus maximowiczii* Schrenk, 1858 (Rodentia, Cricetidae) // 11th International Conference Rodens et Spatiun on Rodent Biology. Myshkin. P.48.
- Korobitsyna K.V., Kartavtseva I.V., Frisman L.V., Kryukov A.P. & Voyta L.L. 2005. [Chromosomal polymorphism and allozyme differentiation in vole *Microtus maximowiczii* Schrenk, 1858 from Transbaikalian region] // [Ecosystems of Mongolia and Frontier Areas of Adjacent Countries: Natural Resources, Biodiversity and Ecological Prospects.] Ulaanbaatar. P.287–289 [in Russian].
- Kovalskaya Yu.M. 1977. [Chromosome polymorphism in *Microtus maximowiczii* Schrenk, 1858 (Rodentia, Cricetidae)] // Byulleten' Moskovskogo Obshchestva Ispytatelei Prirody. Vol.82. No.2. P.38–48 [In Russian, with English summary].
- Kovalskaya Yu.M., Khotolkhu N. & Orlov V.N. 1980. [Geographical distribution of chromosome mutations and structure of the species *Microtus maximowiczii* (Rodentia, Cricetidae)] // Zoologicheskii Zhurnal. Vol.59. No.12. P.1862–1869 [In Russian, with English summary].
- Meyer M.N. 1978. [Taxonomy and intraspecific variation of *Microtus* from the Far East of the USSR (Rodentia,

- Cricetidae)] // Trudy Zoologicheskogo Instituta AN SSSR. Vol.75. P.3–62 [in Russian].
- Meyer M.N., Golenishchev F.N., Radjabli S.I. & Sablina O.V. 1996. [Grey voles (subgenus *Microtus*) of Russia and adjacent territories] // Trudy Zoologicheskogo Instituta RAS. T.232. P.1–320 [in Russian].
- Meyer M.N., Jordan M. & Walknowska J. 1967. A karyosystematic study of some *Microtus* species // Folia Biologica (Krakow). Vol.15. No.3. P.251–264.
- Sheremetyeva I.N., Kartavtseva I.V., Haring E., Frisman L.V. & Kryukov A.P. 2008. Morphological and genetic characteristics of *Microtus maximowiczii gromovi* Vorontsov, Boeskorov, Lyapunova et Revin, 1988 // 11th International Conference Rodens et Spatiun on Rodent Biology. Myshkin. P.84.
- Vorontsov N.N., Boeskorov G.G., Lyapunova E.A., & Revin Yu.V. 1988. [A new chromosome form and variability of molars in voles *Microtus maximowiczii* (Rodentia, Cricetidae)] // Zoologicheskii Zhurnal. Vol.67. No.2. P.205–215 [in Russian].